



# Evenity (romosozumab-aqqg) Injection Orders

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_  Male  Female

Diagnosis (please provide ICD10 code)  Age-related osteoporosis without current pathological fracture M81.0

Age-rel osteopor w current path fracture, unsp site, init. M80.00XA  NKDA Allergies: \_\_\_\_\_

New Start Therapy  Continuation of Therapy Date of last dose (if applicable): \_\_\_\_\_

## Ordering Provider:

Provider NPI: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_

Practice Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip Code: \_\_\_\_\_

### TRIED AND FAILED MEDICATIONS:

- Fosamax  Boniva  Actonel  Evista  Reclast  
 Contraindications to above: \_\_\_\_\_

\*NOTE: As of January 2023 most major Insurance plans require trial/failure of both oral and IV bisphosphonate therapy (such as Reclast and Prolia) before approving Evenity treatment. If your patient has not tried and failed or has a contraindication to oral or IV bisphosphonate and you still would like to pursue Evenity treatment, we encourage an addendum, progress note or letter of medical necessity explaining why step therapy is not recommended/beneficial for the patient i.e contraindication, intolerance, allergy, etc.

### REQUIRED TESTING/LABS:

- Clinical/Progress Notes supporting primary diagnosis (please attach)
- DEXA scan results and date (please attach): \_\_\_\_\_
- Calcium level and date (please attach most recent CMP): \_\_\_\_\_

## EVENITY ORDERS

### DOSING:

- Evenity 210mg (two 105mg prefilled syringes) subcutaneous injections

### FREQUENCY:

- Once a month for 12 doses

### Infuse One Standing Orders:

- Provide treatment under Infuse One's Clinical Guidelines, Medication Safety Protocol, Emergency Guidelines, and Action Plan for Infusion Reactions.

### REFILLS:

- \_\_\_\_\_  
*(if not indicated prescription will expire one year from date signed)*

\_\_\_\_\_  
 Provider Name

\_\_\_\_\_  
 Provider Signature

\_\_\_\_\_  
 Date

Evenity may increase the risk of myocardial infarction, stroke and cardiovascular death. Evenity should not be initiated in patients who have had a myocardial infarction or stroke within the preceding year. Consider whether the benefits outweigh the risks in patients with other cardiovascular risk factors. Instruct patients to take calcium 1000 mg daily and at least 400 IU vitamin D daily. Hypocalcemia must be corrected prior to beginning Evenity.