

Reclast (zoledronic acid) Infusion Orders

Patient Name:	DOB:	[☐ Male ☐ Female
Diagnosis (please provide ICD10 code)			
□ Other:			
☐ New Start Therapy ☐ Continuation of Therapy	Date of last dose (if applicable):	□ NKDA A	Allergies:
Ordering Provider:			
Provider NPI:	Phone:	Fax:	
Practice Address:	City:	State:	Zip Code:
PREMEDICATIONS:		REQUIRED TES	TING/LABS:
☐ Acetaminophen 1000mg PO ☐ Solu-Medro	n izanig ivi	☑ Clinical/Progress Notes supporting primary diagnosis (please attach)	
☐ Diphenhydramine 25mg PO ☐ Solu-Cortef	100mg IVP	DEXA scan result	ts and date (please attach):
☐ Ceterizine 10mg PO ☐ Diphenhydi	ramine 25mg IVP	——————————————————————————————————————	 P lab results (please attach)
☐ Other:		iviost recent civii	ido results (piedse detdeti)
RECLAST ORDERS			
DOSING:	I	REFILLS:	
☑ Reclast 5mg/100ml IV infusion over at leas	t 15 minutes	No Refills	
FREQUENCY:			
☑ Once annually			
Infuse One Standing Orders:			
Provide treatment under Infuse One's Clinical Guand Action Plan for Infusion Reactions.	uidelines, Medication Safety Protocol	Emergency Guide	elines,
Provider Name			
Provider Signature		Date	

Hypocalcemia may worsen during treatment. Patients must be adequately supplemented with calcium and vitamin D. Renal Impairment: Monitor creatinine clearance before each dose.

Osteonecrosis of the Jaw (ONJ) has been reported. All patients should have a routine oral exam by the prescriber prior to treatment. Atypical Femur Fractures have been reported. Patients with thigh or groin pain should be evaluated to rule out a femoral fracture. Severe Bone, Joint, and Muscle Pain may occur. Withhold future doses of Reclast if severe symptoms occur.