



Soliris Infusion Order

Patient Name: _____ DOB: _____ Male Female

Diagnosis (please provide ICD10 code) _____

Other: _____

NKDA Allergies: _____

New Start Therapy Continuation of Therapy Date of last dose (if applicable): _____

Ordering Provider:

Provider NPI: _____ Phone: _____ Fax: _____

Practice Address: _____ City: _____ State: _____ Zip Code: _____

PRE-MEDICATION

- Acetaminophen 1000mg PO Solu-Medrol 125mg IVP
- Diphenhydramine 25mg PO Solu-Cortef 100mg IVP
- Ceterizine 10mg PO Diphenhydramine 25mg IVP

REQUIRED LABS

- Meningococcal vaccination (both conjugate and serogroup B) are required prior to initiating Soliris infusions (please attach documentation).
- Date of meningococcal vaccine: _____

SOLIRIS ORDERS

- Induction Dose (**Choose one. If patient has already completed induction dose, proceed to maintenance dose.**)

- 600mg weekly for the first four weeks followed by 900mg for the fifth dose one week later, then 900mg two weeks later
- 900mg weekly for the first four weeks followed by 1200mg for the fifth dose one week later, then 1200mg two weeks later

- Maintenance Dose (**Choose one**)

- 900mg every two weeks
- 1200mg every two weeks

- Dilute with 0.9% NS to a final concentration of 5mg/ml
(600mg doses final volume 120ml, 900mg doses final volume 180ml, 1200mg doses final volume 240ml)
- Infuse over 35 minutes in adults

Refills: _____ (if not indicated prescription will expire one year from date signed)

Infuse One Standing Orders:

- Provide treatment under Infuse One's Clinical Guidelines, Medication Safety Protocol, Emergency Guidelines, and Action Plan for Infusion Reactions.

Provider Name

Provider Signature

Date

Monitor the patient for at least one hour following completion of the infusion for signs or symptoms of an infusion reaction.

Soliris is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS). Under the Soliris REMS, prescribers must enroll in the program.