## FAX ORDER TO 561-516-6626



Gastroenterology Referral Form					
**Please Attach Copy of Insurance Cards (Front & Back)**					
Last Name: First Name:		Name:	DOB:	Practice:	
Address:				Address:	
City:	State	:: Zip:	Sex: M F	City: State: 2	Zip:
Phone:		SSN#		Prescriber Name:	
Insurance Information Prescriber NPI:					
Insurance Plan: Insur		Insurance Plan:		Nurse/Key Contact:	
Policy #		Policy #		Phone:	
Plan I.D. #		Plan I.D. #		Fax: Email:	
Diagnosis & Clinical Information					
**Please Attach Clinical/Progress Notes, Labs, Test, Supporting Primary Diagnosis**					
Crohn's Disease     Diagnosis code:       Ulcerative Colitis     Diagnosis code:					
	2.43		Allergies:		
Other:					
Currently received and/or prior filed therapies: NKDA					
Length of treatment:			Height: Weight:		
Reason for disco	ontinuation:		Site of Care: 🔄 Home	AIC Other:	
Prescription Information					
		Prescript	tion Information		
Medication	Dose/Strength	Prescript	tion Information Directions		Refills
				y 8 weeks thereafter	Refills
Medication	Dose/Strength	INITIAL: Infuse 300mg	Directions		Refills
Entyvio (vedolizumab)		INITIAL: Infuse 300mg	<b>Directions</b> g IV at week 0, 2, 6, then every	reeks	Refills
Entyvio	300mg vial	INITIAL: Infuse 300mg MAINTENANCE: Infus	Directions g IV at week 0, 2, 6, then every e 300mg IV every w	reeks then every 8 weeks thereafter	Refills
Entyvio (vedolizumab)		INITIAL: Infuse 300mg MAINTENANCE: Infus	Directions g IV at week 0, 2, 6, then every e 300mg IV every w mg/kg IV at week 0, 2, 6,	reeks then every 8 weeks thereafter	Refills
Entyvio (vedolizumab)	300mg vial	INITIAL: Infuse 300mg MAINTENANCE: Infuse INITIAL: Infuse MAINTENANCE: Infuse Other	Directions g IV at week 0, 2, 6, then every e 300mg IV every w mg/kg IV at week 0, 2, 6, e mg/kg IV every	reeks then every 8 weeks thereafter	Refills
Entyvio (vedolizumab)	300mg vial	INITIAL: Infuse 300mg MAINTENANCE: Infuse INITIAL: Infuse MAINTENANCE: Infuse Other Pharmacist will round	Directions           g IV at week 0, 2, 6, then every           e 300mg IV everyw          mg/kg IV at week 0, 2, 6,           emg/kg IV every           It to the nearest 100mgGi	veeks then every 8 weeks thereafter weeks	Refills
Entyvio (vedolizumab)	300mg vial	<ul> <li>INITIAL: Infuse 300mg</li> <li>MAINTENANCE: Infuse</li> <li>INITIAL: Infuse</li> <li>MAINTENANCE: Infuse</li> <li>Other</li> <li>Pharmacist will round</li> <li>INITIAL: Weight based</li> <li>55kg or less: 2</li> </ul>	Directions g IV at week 0, 2, 6, then every e 300mg IV every wmg/kg IV at week 0, 2, 6, emg/kg IV every I to the nearest 100mgGi dosing, infuse IV 60mg (2 vials)5	veeks then every 8 weeks thereafter weeks	Refills
Entyvio (vedolizumab)	300mg vial	INITIAL: Infuse 300mg MAINTENANCE: Infuse INITIAL: Infuse MAINTENANCE: Infuse MAINTENANCE: Infuse Other Pharmacist will round INITIAL: Weight based 55kg or less: 2 Greater than 88	Directions g IV at week 0, 2, 6, then every e 300mg IV every wmg/kg IV at week 0, 2, 6, emg/kg IV every I to the nearest 100mgGi dosing, infuse IV 60mg (2 vials)5 5kg: 520 mg (4 vials)	veeks then every 8 weeks thereafter weeks ve exact dose (do NOT round) 5kg to 85kg: 390mg (3 vials)	Refills
Entyvio (vedolizumab)	300mg vial	INITIAL: Infuse 300mg MAINTENANCE: Infuse INITIAL: Infuse MAINTENANCE: Infuse MAINTENANCE: Infuse Other Pharmacist will round INITIAL: Weight based 55kg or less: 2 Greater than 88	Directions g IV at week 0, 2, 6, then every e 300mg IV every wmg/kg IV at week 0, 2, 6, emg/kg IV every I to the nearest 100mgGi dosing, infuse IV 60mg (2 vials)5 5kg: 520 mg (4 vials)	veeks then every 8 weeks thereafter weeks ve exact dose (do NOT round)	Refills
Entyvio (vedolizumab)  Inflectra (infliximab)  Remicade  Renflexis  Stelara (ustekinumab)	300mg vial 100mg vial 130 mg / 26ml vial 90mg (2x 45mg vials)	<ul> <li>INITIAL: Infuse 300mg</li> <li>MAINTENANCE: Infuse</li> <li>INITIAL: Infuse</li> <li>MAINTENANCE: Infuse</li> <li>Other</li> <li>Other</li> <li>Pharmacist will round</li> <li>INITIAL: Weight based</li> <li>55kg or less: 2</li> <li>Greater than 88</li> <li>MAINTENANCE: Inject</li> </ul>	Directions g IV at week 0, 2, 6, then every e 300mg IV every wmg/kg IV at week 0, 2, 6, emg/kg IV every I to the nearest 100mgGi dosing, infuse IV 60mg (2 vials)5 5kg: 520 mg (4 vials)	veeks then every 8 weeks thereafter weeks ve exact dose (do NOT round) 5kg to 85kg: 390mg (3 vials)	Refills
Entyvio (vedolizumab)	300mg vial	<ul> <li>INITIAL: Infuse 300mg</li> <li>MAINTENANCE: Infuse</li> <li>INITIAL: Infuse</li> <li>MAINTENANCE: Infuse</li> <li>Other</li> <li>Pharmacist will round</li> <li>INITIAL: Weight based</li> <li>55kg or less: 2</li> <li>Greater than 84</li> <li>MAINTENANCE: Inject</li> <li>INITIAL: Infuse 600mg</li> </ul>	Directions g IV at week 0, 2, 6, then every e 300mg IV every w mg/kg IV at week 0, 2, 6, e mg/kg IV every I to the nearest 100mg Gi dosing, infuse IV 60mg (2 vials) 5 5kg: 520 mg (4 vials) 90mg SQ 8 weeks after initial y/10mL IV at week 0, 4, and 8	veeks then every 8 weeks thereafter weeks ve exact dose (do NOT round) 5kg to 85kg: 390mg (3 vials)	
	300mg vial 100mg vial 130 mg / 26ml vial 90mg (2x 45mg vials)	<ul> <li>INITIAL: Infuse 300mg</li> <li>MAINTENANCE: Infuse</li> <li>INITIAL: Infuse</li> <li>MAINTENANCE: Infuse</li> <li>Other</li> <li>Pharmacist will round</li> <li>INITIAL: Weight based</li> <li>55kg or less: 2</li> <li>Greater than 88</li> <li>MAINTENANCE: Inject</li> <li>INITIAL: Infuse 600mg</li> <li>MAINTENANCE: Inject</li> </ul>	Directions g IV at week 0, 2, 6, then every e 300mg IV every w mg/kg IV at week 0, 2, 6, e mg/kg IV every I to the nearest 100mg Gi dosing, infuse IV 60mg (2 vials) 5 5kg: 520 mg (4 vials) 90mg SQ 8 weeks after initial //10mL IV at week 0, 4, and 8 360mg/2.4ml SQ via injector	reeks then every 8 weeks thereafter 	
	300mg vial         100mg vial         130 mg / 26ml vial         90mg (2x 45mg vials)         600mg / 10 ml vial	INITIAL: Infuse 300mg MAINTENANCE: Infuse INITIAL: Infuse MAINTENANCE: Infuse Other Other Pharmacist will round INITIAL: Weight based 55kg or less: 24 Greater than 88 MAINTENANCE: Inject INITIAL: Infuse 600mg MAINTENANCE: Inject Acetaminophen	Directions g IV at week 0, 2, 6, then every e 300mg IV every w mg/kg IV at week 0, 2, 6, e mg/kg IV every I to the nearest 100mg Gi dosing, infuse IV 60mg (2 vials) 5 5kg: 520 mg (4 vials) 90mg SQ 8 weeks after initial y/10mL IV at week 0, 4, and 8	reeks then every 8 weeks thereafter 	
	300mg vial         100mg vial         130 mg / 26ml vial         90mg (2x 45mg vials)         600mg / 10 ml vial         • other medications	<ul> <li>INITIAL: Infuse 300mg</li> <li>MAINTENANCE: Infuse</li> <li>INITIAL: Infuse</li> <li>MAINTENANCE: Infuse</li> <li>Other</li> <li>Pharmacist will round</li> <li>INITIAL: Weight based</li> <li>55kg or less: 21</li> <li>Greater than 88</li> <li>MAINTENANCE: Inject</li> <li>INITIAL: Infuse 600mg</li> <li>MAINTENANCE: Inject</li> <li>Acetaminophen</li> <li>Diphenhydramine</li> </ul>	Directions g IV at week 0, 2, 6, then every e 300mg IV everywmg/kg IV at week 0, 2, 6, emg/kg IV every I to the nearest 100mgGi dosing, infuse IV 60mg (2 vials)5 5kg: 520 mg (4 vials) 90mg SQ 8 weeks after initial p/10mL IV at week 0, 4, and 8 360mg/2.4ml SQ via injector mg PO prior to infusior mg POIV	then every 8 weeks thereafter weeks we exact dose (do NOT round) 5kg to 85kg: 390mg (3 vials) dose, then every 8 weeks thereafter at week 12, then every 8 weeks thereaft	
	300mg vial         100mg vial         130 mg / 26ml vial         90mg (2x 45mg vials)         600mg / 10 ml vial         600mg / 10 ml vial	<ul> <li>INITIAL: Infuse 300mg</li> <li>MAINTENANCE: Infuse</li> <li>INITIAL: Infuse</li> <li>MAINTENANCE: Infuse</li> <li>Other</li> <li>Pharmacist will round</li> <li>INITIAL: Weight based</li> <li>55kg or less: 2i</li> <li>Greater than 89</li> <li>MAINTENANCE: Inject</li> <li>INITIAL: Infuse 600mg</li> <li>MAINTENANCE: Inject</li> <li>Acetaminophen</li> <li>Diphenhydramine</li> <li>250ml 0.9%NaCl for hy</li> </ul>	Directions g IV at week 0, 2, 6, then every e 300mg IV everywmg/kg IV at week 0, 2, 6, emg/kg IV every I to the nearest 100mgGi dosing, infuse IV 60mg (2 vials)5 5kg: 520 mg (4 vials) 90mg SQ 8 weeks after initial p/10mL IV at week 0, 4, and 8 360mg/2.4ml SQ via injector mg PO prior to infusior mg POIV	veeks then every 8 weeks thereafterweeks ve exact dose (do NOT round) 5kg to 85kg: 390mg (3 vials) dose, then every 8 weeks thereafter at week 12, then every 8 weeks thereaft <b>Flush Protocol</b> * NaCl 0.9% 10ml	
	300mg vial         100mg vial         130 mg / 26ml vial         90mg (2x 45mg vials)         600mg / 10 ml vial         600mg / 10 ml vial	<ul> <li>INITIAL: Infuse 300mg</li> <li>MAINTENANCE: Infuse</li> <li>INITIAL: Infuse</li> <li>MAINTENANCE: Infuse</li> <li>Other</li> <li>Pharmacist will round</li> <li>INITIAL: Weight based</li> <li>55kg or less: 21</li> <li>Greater than 88</li> <li>MAINTENANCE: Inject</li> <li>INITIAL: Infuse 600mg</li> <li>MAINTENANCE: Inject</li> <li>Acetaminophen</li> <li>Diphenhydramine</li> </ul>	Directions g IV at week 0, 2, 6, then every e 300mg IV everywmg/kg IV at week 0, 2, 6, emg/kg IV every I to the nearest 100mgGi dosing, infuse IV 60mg (2 vials)5 5kg: 520 mg (4 vials) 90mg SQ 8 weeks after initial p/10mL IV at week 0, 4, and 8 360mg/2.4ml SQ via injector mg PO prior to infusior mg POIV	veeks then every 8 weeks thereafterweeks ve exact dose (do NOT round) 5kg to 85kg: 390mg (3 vials) dose, then every 8 weeks thereafter at week 12, then every 8 weeks thereaft <b>Flush Protocol</b> * NaCl 0.9% 10ml	
	300mg vial     300mg vial     100mg vial     130 mg / 26ml vial     90mg (2x 45mg vials)     600mg / 10 ml vial     600mg / 10 ml vial	INITIAL: Infuse 300mg MAINTENANCE: Infuse INITIAL: Infuse MAINTENANCE: Infuse Other Pharmacist will round INITIAL: Weight based 55kg or less: 2 Greater than 89 MAINTENANCE: Inject INITIAL: Infuse 600mg MAINTENANCE: Inject Acetaminophen Diphenhydramine 250ml 0.9%NaCl for hy Other	Directions g IV at week 0, 2, 6, then every e 300mg IV everywmg/kg IV at week 0, 2, 6, emg/kg IV every I to the nearest 100mgGi dosing, infuse IV 60mg (2 vials)5 5kg: 520 mg (4 vials) 90mg SQ 8 weeks after initial //10mL IV at week 0, 4, and 8 360mg/2.4ml SQ via injector mg PO prior to infusior mg PO prior to infusior mg POIV //dration	veeks then every 8 weeks thereafterweeks ve exact dose (do NOT round) 5kg to 85kg: 390mg (3 vials) dose, then every 8 weeks thereafter at week 12, then every 8 weeks thereaft <b>Flush Protocol</b> * NaCl 0.9% 10ml	
	300mg vial     300mg vial     100mg vial     130 mg / 26ml vial     90mg (2x 45mg vials)     600mg / 10 ml vial     600mg / 10 ml vial     so ther medications lies as per protocol cit as per protocol cit as per protocol cit as per protocol	<ul> <li>INITIAL: Infuse 300mg</li> <li>MAINTENANCE: Infuse</li> <li>INITIAL: Infuse</li> <li>MAINTENANCE: Infuse</li> <li>Other</li> <li>Pharmacist will round</li> <li>INITIAL: Weight based</li> <li>55kg or less: 2</li> <li>Greater than 84</li> <li>MAINTENANCE: Inject</li> <li>INITIAL: Infuse 600mg</li> <li>MAINTENANCE: Inject</li> <li>INITIAL: Infuse 600mg</li> <li>MAINTENANCE: Inject</li> <li>Acetaminophen</li> <li>Diphenhydramine</li> <li>250ml 0.9%NaCl for hy</li> <li>Other</li> </ul>	Directions g IV at week 0, 2, 6, then every e 300mg IV everywmg/kg IV at week 0, 2, 6, emg/kg IV at week 0, 2, 6, emg/kg IV every I to the nearest 100mgGi dosing, infuse IV 60mg (2 vials)5 5kg: 520 mg (4 vials) 90mg SQ 8 weeks after initial y/10mL IV at week 0, 4, and 8 360mg/2.4ml SQ via injector mg PO prior to infusior mg POIV ydration	then every 8 weeks thereafter weeks we exact dose (do NOT round) 5kg to 85kg: 390mg (3 vials) dose, then every 8 weeks thereafter at week 12, then every 8 weeks thereaft <b>Flush Protocol</b> * NaCl 0.9% 10ml * Before & after infusion	

PRESCRIBER MUST MANUALLY SIGN - STAMP SIGNATURE, SIGNATURE BY OTHER PERSONNEL AND COMPUTER-GENERATED SIGNATURES WILL NOT BE ACCEPTED The attached document(s) contain confidential information which may be considered to be Protected Health Information and therefore required to be maintained as private and secure under HIPAA. The documents may also contain information which is otherwise considered to be privileged under state and federal laws. This communication is for the intended recipient only. If you are not the intended recipient, or a person responsible for delivering this communication to the intended recipient, you are prohibited from viewing, copying and/or distributing the information contained herein. Unlawful disclosure of the information attached may subject you to monetary penalties and sanctions. If you have received this communication in error, you should notify the sender immediately and thereafter permanently destroy all copies of this document in its entirety.

This form is not considered an order or prescription for medical services and/or supplies unless and until it is formally authorized by a healthcare provider in compliance with applicable laws and regulations.