

NULOJIX (belatacept)

| PATIENT INFORMATION | | |
|---|---|---|
| Patient Name: Status: New to therapy Continuing Next Due Date (if applicable PROVIDER INFORMATION | Date of Birth: | Phone: |
| | | |
| Provider Name: | Pi | rovider NPI: |
| Practice Address: | City: | State: Zip: |
| Practice Name: | | |
| Practice Phone: Fax: | Contact Pe | erson: |
| MEDICAL INFORMATION | | |
| Patient Weight: Patient Height: ICD-10 Code (required | uired): ICD-10 Description: | |
| Known Allergies: | | |
| Required Labs: EBV serostatus and TB screening | | |
| DETAILS NEEDED FOR AUTHORIZATION | | |
| Proof of patient being concurrently treated with any other biologic: | | |
| Nolojix Distribution Program Notified (855) 511-6180 - Patient ID | | |
| NULOJIX (belatacept) ORDERS | | |
| Dosing for Initial Phase and Initial Maintenance mg IV on Day 1 (day of transplantation, prior to transplantation) and Day 5 (approx. 96 hours after Day 1 dose), at the end of week 2, week 4, week 8, and week 12 after transplantation. Thenmg IV at the end of week 16 after transplantation and every 4 weeks (plus or minus 3 days) thereafter. * Patient has received doses thus far, next dose due on | * Prescribed dose must be evenly divisible by 12.5mg * The total infusion dose of Nulojix should be based on the actual body weight of the patient at the time of transplantation, and should not be modified during the course of therapy, unless there is a change in the body weight of greater than 10%. If the patient has had >10% weight change, please notify the physician for dose change recommendations. Additional Orders/Comments: | |
| Dosing for Maintenance Phase | | |
| ADULT RESCUE MANAGEMENT PROTOCOL | | |
| These include fever, chills, rigors, headache, rash, itching, swelling, edema, nausea, vomiting, abdominal pain, hypotension, and respiratory distress. Following standing reaction orders, including diphenhydramine, methylprednisolone, albuterol, and oxygen as needed. For severe reactions, administer Epi-pen or equivalent and call 911. Repeat if severe symptoms persist. REQUIRED STANDARD DOCUMENTATION NEEDED | | |
| | | if available, lact history and physical |
| Patient demographics Patient medical insurance card, copied front and back Patient pharmacy card, copied front and back (if they have one) | Most recent chart notes and if available, last history and physical. All relevant scans, tests and laboratory results. If new medication for patient, chart notes which include decision to begin treatment. If not, provide last treatment date. | |
| PROVIDER AUTHORIZATION | | |
| | | |
| Provider's Signature: | Print Name: | Date: |