

Email to: intake@infuseone.com Fax to: 888-261-6644

]	Palm	Beach	Gardens

- ☐ Melbourne
- □ Tallahassee
- ☐ Fort Lauderdale

Dermate	ology
Referral	Form

_ast Name, First Name:]	Date of Birth: Gender: \square M \square F \square Other		ner	
ddress:		(City, State, Zip:			
hone:			SSN#			
eferring Practice:						
ractice Address:		(City, State, Zip:			
Prescriber Name:		F	Prescriber NPI:			
urse/Key Contact	•	F	Phone:			
IX:			Email:			
Nursino	g & Lab Orders					
urse Orders: Nur	se to provide assessment, teaching,	n and as needed <i>Heparin</i> - 🗆 10units/mL -		ice insertion and/or management per physician of 5mL flush after post-infusion NS flush if indicated to ma		
Prescri	otion Orders					
Anaphylaxis Kit: (Check all that apply)	☐ Epinephrine 0.3mg IM as needed ☐ Diphenhydraminemg IV infus ☐ Acetaminophenmg PO	sion as needed NS Hydration 500 minutes prior to infusion Solu-1	ml IV infusion over 30 minu Medrolmg IVminute	es prior to infusion		
Check all that apply) Supply Orders: All	☐ Diphenhydramine mg ☐ Posupplies for vascular access line care, dru					
			will be provided as necessar	ry		
	otion Information					
PRODUCT			CTIONS		REFILLS	
	? Yes no if no, when was la		When is patient due	for next dose?		
☐ Ilumya ☐ Infliximab ☐ Avsola	100mg sc injection at 0 and 4 weeks then every 12 weeks INDUCTION:mg/kg ormg iv infusion via gravityOR pump over at least 2 hours at weeks 0, 2, and 6					
☐ Inflectra ☐ Remicade ☐ Renflexis ☐ Renflexis ☐ Inflectra ☐ MAINTENANCE:mg/kg mg iv infusion via ☐ grave (Note: round to nearest 100mg for medicaid patients) ☐ If remicade infusion tolerated, adjust infusion time according to ma			acturer package insert.			
☐ Simponi Aria	2 mg/kg IV infusion via □ gravity	OR pump over 30 minutes	s at weeks 0 and 4, and e	very 8 weeks thereafter		
☐ Spevigo	☐ 900 Mg iv infusion over 90 min	utes 🔲 Additional 900 mg iv infus	ion over 90 minutes one	week after initial dose if flare symptoms persist		
⊐ Stelara	☐ For patients > 100 kg, 90 mg S Psoriasis Pediatric Patients 6 to ☐ For patients <= 60 kg, 0.75 mg ☐ For patients 60 kg — 100kg, 45 ☐ For patients >100kg, 90 mg SC Psoriasis Adult Subcutaneous	SC injection initially and 4 weeks later to injection initially and 4 weeks later 17 (based on weight at time of in /kg SC injection initially and 4 weeks mg SC injection initially and 4 weeks later, followed by 45 mg SC weeks later.	r, followed by 90 mg even itial dose) s later, then every 12 week is later, then every 12 week then every 12 weeks	y 12 weeks ks ks		
	☐ 45 mg SC injection initially and 4 weeks later, followed by 45 mg SC injection every 12 weeks ☐ For patients with co-existent moderate-to-severe plaque psoriasis weighing >100 kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks					
Xolair	☐ 150 or ☐ 300 mg SC injection once every 4 weeks					
☐ IG ☐ Other	For Immunoglobulin therapy pled	se refer to Immunoglobulin Form				
signing this for escription insures	ance companies. RITTEN			tion designated agent in dealing with medica		
		Print Name: _		Date:		
GUBSTITUTION PERMITTED Prescriber's Signature: P				Nate:		
	o attach the following:	Provider Phone Line: 833–88		Date		
☐ Patient demog	graphics & front/back copy of all insu visit notes, history & physical, lab & p ation list & list of prior medications t	ertinent procedure results	☐ HBV lab results with	last 12 months (Stelara, Simponi Aria, Ilumya & Inf in last 12 months (Infliximabs & Simponi Aria only) essity if drug dosing or indication is outside of FD/		