



TzielD® (teplizumab-mzwv) Plan of Treatment

PATIENT DEMOGRAPHICS:

Patient Name:		Patient's Phone:	
Date of Birth:		Address:	
Allergies:		City, State, Zip:	
Height in inches:	Weight: LB or KG	Gender:	Allergies: <input type="checkbox"/> See list <input type="checkbox"/> NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

E10.9 - Type 1 diabetes mellitus without complications	-Other:
E10.8 - Type 1 diabetes mellitus with unspecified complications	

REQUESTED DOCUMENTATION:

REQUESTED DOCUMENTATION:		HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE? Yes or No	
1	Insurance information	PLEASE STATE REQUIRED WASHOUT FROM PREVIOUS THERAPY:	Continue current order until insurance approved
2	H&P including tried and failed therapies		
3	Full medication list		
4	Required: Recent CBC with diff and LFTs		

HOME SUPPLY ORDER: *Applicable to all home infusion patients subject to insurance approval

All supplies for vascular access line care, dressing kit, drug administration, adverse reaction kit, Infusion pump, IV pole, pole clamp etc. will be

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive teplizumab-mzwv if receiving antibiotics for active infectious process, active fever and/or suspected infection or surgery. If patient experiences severe cytokine release syndrome, consider temporarily pausing therapy for 1-2 days and review prescribing information before continuing therapy.

Home Anaphylaxis Kit: Dispense and administer for mild and severe reaction. Applicable to all home infusion patients.

2 - Epinephrine 1 mg/ml 1 ml
 2 - Diphenhydramine 50 mg/ml 2 ml vial
 Syringes, needles and 0.9% Normal Saline Flushes 10 mls to administer
 Complete Home Infusion Physician Standing Order for complete Home Infusion, Nursing, and Adverse Reaction Orders to be utilized in the event of an adverse reaction/anaphylaxis.
 Epinephrine administered IM per weight based dosing guide and Benadryl 25-50mg IVP to be administered by clinician in the home.

***Per FDA labeling: Premedication of Acetaminophen or NSAID and antihistamine, and/or an antiemetic is suggested for at least the first 5 days of the 14-day treatment course**

***Any selected premedication will only be given with infusions 1 through 5 unless otherwise designated here:**

IV	PO						
	Diphenhydramine	25mg	50mg	325mg	500mg	650mg	1000mg
Methylprednisolone	40mg	125mg	Famotidine	20mg	40mg		
Ondansetron	4mg	8mg	Diphenhydramine	25mg	50mg		
Other:			Fexofenadine	60mg	180mg		
			Cetirizine	10mg			
			Loratadine	10mg			
			Other:				

DOSAGE:

Treatment Day	Dosage
Day 1	65 mcg/m2
Day 2	125 mcg/m2
Day 3	250 mcg/m2
Day 4	500 mcg/m2
Day 5-14	1030 mcg/m2

LAB ORDERS:

Draw CBC with diff & LFT on infusion days 5 and 8

Other: _____

MEDICATION:

TzielD® dose to be diluted in 25ml NS administered as an IV infusion over a minimum of 30-minutes. Administer once daily for 14 consecutive days.

Follow first 5 infusions with a 1-hour post infusion observation.

Physician responsible for all follow up lab monitoring.

Per FDA labeling, with AST/ALT > 5x ULN, bilirubin > 3x ULN, or Lymphocyte count <500/mcL lasting one week or longer, patient may be ineligible to receive TzielD®

SPECIAL/OTHER LAB ORDERS:

SITE OF CARE:

AIC AIC and Home with a nurse

**If site of care not indicated, PIS will coordinate with patient*

LINE USE/CARE ORDERS:

<input checked="" type="checkbox"/> Start PIV/Access CVC <input checked="" type="checkbox"/> Flush PIV/Access per PIV/PICC/CVC protocol. <input checked="" type="checkbox"/> Provide nursing care per Infuse One's Infusion Nursing Procedures and post procedure observation if indicated.	Dispense and Administer as Prescribed
---	--

PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE:

Dispense as written/Brand medically necessary	Substitution permitted